



GrazonPD3™ Herbicide

GrazonPD3™ herbicide provides solid general weed control with residual properties at exceptional value per acre. It has broad use rates and has the flexibility to be a good early season option.

- A higher load formulation of Grazon® P+D herbicide.
- Offers better residual than 2,4-D or dicamba alone.
- Choline formulated – meaning near zero volatility.
- Labeled for aerial and ground broadcast treatments.

Fallow Cropland Opportunity

For control of broadleaf annual and perennial weeds and certain woody plants not rotated to a broadleaf crop.

Rates and Recommendations

Applying GrazonPD3 alone or in tank mix combination with other registered herbicides can be a weed control solution in fallow cropland.

Apply in 2 or more gallons of water per acre by air or 5 or more gallons per acre by ground.

Do not apply more than 0.25 lb ae picloram (2.5 pint of GrazonPD3) per acre as a broadcast treatment per annual growing season.

For best weed control results, do not cultivate for at least 2 weeks after application or until top growth is dead.

Preplant Interval to Small Grains

- For applications up to 1 pint per acre, allow a minimum of 45 days of soil temperature above 40°F between application and planting.
- For applications greater than 1 pint per acre, allow a minimum of 60 days of soil temperature above 40°F between application and planting.
 - »» 90 days for North Dakota, Nebraska, Montana, South Dakota and Wyoming.

Crop Rotation


Use only on land to be planted the following year to grass, barley, oats, wheat, forage sorghum, sudangrass, or fallowed.

- Do not plant grain sorghum within 8 months after application.
- Do not use this product for sweet sorghum production or on land that will be rotated to sweet sorghum.

Many broadleaf crops are extremely sensitive to soil residues of GrazonPD3 herbicide. Do not plant sensitive broadleaf crops for 36 months after treatment or until soil residues have declined to a safe level as indicated by an adequately sensitive bioassay using the intended broadleaf crop.

Conduct a bioassay following treatment and prior to planting any sensitive broadleaf crop.

- Do not rotate to forage sorghum if greater than 2 pints per acre of GrazonPD3 has been applied.
- Do not plant forage sorghum within 8 months after application.
- Do not use this product for sweet sorghum production or on land that will be rotated to sweet sorghum.



Russian thistle



Pigweed

Weed Control Recommendations

Annual Weeds (such as Russian thistle and wild buckwheat)

Apply .5 to 1 pint per acre of GrazonPD3™ herbicide in tank mix combination with other registered herbicides applying when weeds are actively growing.

Field Bindweed

Apply 1 to 2.25 pints per acre of GrazonPD3 when bindweed is actively growing. Optimum time for treatment is when plant runners reach at least 8 to 12 inches. Use 1 pint per acre to control light to moderate infestations under good growing conditions or to reduce the potential for crop injury. Use 2.25 pint per acre for heavy infestations and to start a treatment program for long-term control. Some regrowth will occur the following season and a re-treatment program of 1 pint of GrazonPD3 for one to two years will provide bindweed stand reduction.

Canada thistle




Apply 2.25 pint per acre of GrazonPD3 when the majority of thistle plants are emerged but prior to bud stage.

Application Recommendations

- Common use rate is 20 fluid ounces per acre for general broadleaf weed control.
- To enhance foliage wetting and coverage, an approved non-ionic surfactant (NIS) may be added to GrazonPD3.

Key Weeds Controlled

Control of Annual and Perennial Broadleaf Weeds	
Annual pricklypoppy	Wild buckwheat
Wild Carrot	Cocklebur
Lambsquarter	Marestail
Morningglory	Prickly lettuce
Pigweed	Puncturevine
Russian thistle	Sunflower

-  Follow us on Facebook @CortevaPastures
-  Follow us on Twitter @CortevaPastures
-  Follow us on Instagram @CortevaPastures



To find your local Corteva Range and Pasture Specialist
 visit rangeandpasture.com/specialist

Visit us at rangeandpasture.com

™ Trademarks of Corteva Agriscience and its affiliated companies.
 GrazonPD3™ and Grazon® P+D are a Restricted Use Pesticide.

This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use, and product warranty and liability limitations, which must be read before using the product(s). Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label direction and precautions for use when using any pesticide alone or in tank-mix combinations.

Contact your state pesticide regulatory agency to determine if a product is registered for sale or use in your state.

Always read and follow label directions. ©2023 Corteva. TSK-10454 COR (03/23)

