

New Era of Burndown Control

HERBICIDE


Active Ingredient

Elevore® herbicide contains Arylex™ Active, an active ingredient from the arylpicolinates family discovered by Corteva Agriscience.

Mode of action	Site of action
Growth regulator	Group 4

Prevents regrowth of emerged weeds for clean fields at planting

Elevore provides thorough control of labeled weeds like henbit, lambsquarters, cutleaf evening-primrose and marestail up to 8" tall by rapidly absorbing into the plant's cells.




Tank-mix compatible with commonly used burndown and residual partners

Elevore can be tank-mixed with a pre-emerge herbicide to start your fields clean at planting and sets the stage for optimum post-emergent applications.

New resistance management tool for ALS and glyphosate resistant weeds

- Low use rate of 1 fl oz./A
- Excellent fit for no-till or reduced tillage production systems
- Labeled for use in soybean, corn and cotton

For more information about Elevore herbicide visit **ElevateYourBurndown.com**, call 800-258-3033 or contact your local Corteva Agriscience territory manager.



Crops	Rotation Interval ¹	
corn	3 ² or 14 days	
barley, canola, millet, oat, rice, rye grasses, sorghum, soybean, sugarcane, sunflower, triticale and wheat (including spring, winter and durum)	14 days	
cotton	30 days	
alfalfa, brassica (cole) leafy vegetables, camelina, chickpea, clover, dry bean, peas (dry and succulent), flax, mustard, peanut, and safflower	9 months	
other crops not listed	15 months	

¹Minimum time that must pass before planting crops after application of Elevore. ²If corn is planted at least 1.5 inches deep and the seed furrow is completely covered with no seed exposed, then planting may occur a **minimum of 3 days following application**.



Visit us at corteva.us

®.TMTrademark of Corteva Agriscience and its affiliated companies.

