

Control pre- and postemergence grass and broadleaf weeds in corn

Eliminate early season competition from grasses and smallseeded broadleaf weeds with soil-applied FulTime® NXT herbicide. Whether used in conventional, reduced-till or no-till production systems, FulTime NXT provides consistent residual control of annual grasses and small-seeded broadleaf weeds


Consistent benefits-whatever the needs are

- Excellent control of annual grasses and broader-spectrum control of broadleaf weeds.
- Contains atrazine for burndown of small weeds at application.
- Requires only 1/4 inch of rain for activation.
- Can be applied from early preplant to early postemergence.
- Time release formulation designed for high residue corn production systems.
- Can be applied alone or in combination with many pre- and postemergence corn herbicides.

Target application timing

Preemergence

Preplant



Use rates (qt./A) by soil texture in conventional, reduced and no-till systems

Soil Texture	Broadcast Rate per Acre (Quarts)*
Coarse	2.9 qt.
Medium	2.9 – 3.7 qt.
Fine	3.2 - 3.7 gt.

*In areas of heavy weed infestation, use up to 4.4 quarts per acre on medium- and fine-textured soils.

For more information about FulTimeNXT herbicide, visit FulTimeNXT.com, call 800-258-3033 or contact your local Corteva Agriscience territory manager.

Receive a broad spectrum of control


FulTime NXT combines the power of acetochlor and atrazine for consistent control of annual grasses and small-seeded broadleaf weeds. See product label for application rates and directions.

Weeds controlled or partially controlled		
Grasses & Sedges		
Barnyardgrass	Oats, wild	
Crabgrass spp.	Panicum, browntop	
Cupgrass, woolly ¹	Panicum, fall	
Foxtail, giant	Panicum, Texas*2	
Foxtail, green	Rice, red	
Foxtail, robust (purple, white)	Sandbur, field*2	
Foxtail, yellow	Shattercane*2	
Goosegrass	Signalgrass, broadleaf	
Johnsongrass, seedling*2	Sprangletop, red	
Millet, wild proso*2	Wheat, volunteer	
Nutsedge, yellow ³	Witchgrass	
Broadleaves		
Beggarweed, Florida	Nightshade, hairy	
Carpetweed	Pigweed, spp.	
Cocklebur ⁴	Purslane, common	
Galinsoga	Pusley, Florida	
Groundcherry, annual	Ragweed, common	
Groundcherry, cutleaf	Ragweed, giant*	
Henbit	Sicklepod*	
Jimsonweed	Sida, prickly	
Kochia ⁵	Smartweed spp.	
Kochia ⁵ Lambsquarters, common	Smartweed spp. Sunflower, common*	
Lambsquarters, common	Sunflower, common*	

Nightshade, black

NOTE: For hard-to-control weeds, additional amounts of Surpass NXT (in corn only) and/or atrazine may be added to the specified treatment rates for

Visit us at corteva.us



®.TMTrademark of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.



¹ Apply 4.4 quarts of FulTime NXT per acre to control this weed in corn; apply 3.7 quarts per acre to control this weed in grain sorghum. Control of this weed can be erratic, especially under dry conditions. Control escaped weeds with cultivation or application of an appropriate registered postemeraence herbicide.

 $^{^{\}rm 2}$ When applied immediately after planting or within 5 days of last tillage, FulTime NXT broadcast applied at a rate of 3.6 to 4.4 quarts per acre in corn will reduce competition from these weeds.

³ Preplant incorporate for control.

⁴ Use the higher rate in the application rate range within each application rate table. Control of these weeds can be erratic, especially under dry weather conditions. Control escaped weeds with cultivation or application of an appropriate registered postemergence herbicide.

⁵ Triazine-resistant biotypes may require a postemergence sequential application of a non-triazine herbicide for control.